


Epidemiology of HIV Infection through 2014

National Center for HIV/AIDS, Viral Hepatitis, STD & TB Prevention
Division of HIV/AIDS Prevention



Diagnoses of HIV Infection among Adults and Adolescents, by Sex, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2014—United States and 6 Dependent Areas

N = 44,609




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Transmission Category, 2014—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU




- Heterosexual contact^a
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnoses of HIV Infection among Adults and Adolescents, by Race/Ethnicity, 2010–2014—United States and 6 Dependent Areas



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Race/Ethnicity, 2014—United States and 6 Dependent Areas



■ American Indian/Alaska Native
■ Asian
■ Black/African American


■ Hispanic/Latino^a
■ Native Hawaiian/other Pacific Islander
■ White

■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^aHispanics/Latinos can be of any race.


Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Age at Diagnosis, 2010–2014—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Diagnoses of HIV Infection among Adults and Adolescents, by Age at Diagnosis, 2014—United States

N = 43,899



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Diagnoses of HIV Infection among Adults and Adolescents, by Transmission Category, 2014—United States and 6 Dependent Areas

Transmission Category	No.	%
Male-to-male sexual contact	29,771	66.7
Injection drug use (IDU)	2,732	6.1
Male-to-male sexual contact and IDU	1,222	2.7
Heterosexual contact ^a	10,781	24.2
Other ^b	103	0.2
Total^c	44,609	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact, by Race/Ethnicity, 2014—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	142	0.5
Asian	799	2.7
Black/African American	11,207	37.6
Hispanic/Latino ^a	7,893	26.5
Native Hawaiian/other Pacific Islander	45	0.2
White	9,012	30.3
Multiple races	674	2.3
Total^b	29,771	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnosed HIV Infections Attributed to Heterosexual Contact^a, by Sex and Race/Ethnicity, 2014—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	10	0.3	36	0.5	46	0.4
Asian	60	1.8	129	1.8	189	1.8
Black/African American	2,113	61.9	4,657	63.2	6,770	62.8
Hispanic/Latino ^b	744	21.8	1,282	17.4	2,026	18.8
Native Hawaiian/other Pacific Islander	1	0.03	7	0.1	8	0.1
White	422	12.4	1,115	15.1	1,537	14.3
Multiple races	63	1.8	142	1.9	205	1.9
Total^c	3,412	100	7,369	100	10,781	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnosed HIV Infections Attributed to Male-to-Male Sexual Contact and Injection Drug Use, by Race/Ethnicity, 2014—United States and 6 Dependent Areas

Race/ethnicity	No.	%
American Indian/Alaska Native	7	0.6
Asian	17	1.4
Black/African American	289	23.6
Hispanic/Latino ^a	257	21.0
Native Hawaiian/other Pacific Islander	1	0.1
White	611	50.0
Multiple races	40	3.3
Total^b	1,222	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnosed HIV Infections Attributed to Injection Drug Use, by Sex and Race/Ethnicity, 2014—United States and 6 Dependent Areas

Race/ethnicity	Males		Females		Total	
	No.	%	No.	%	No.	%
American Indian/Alaska Native	11	0.7	13	1.2	24	0.9
Asian	20	1.2	7	0.7	27	1.0
Black/African American	682	40.8	451	42.5	1,133	41.5
Hispanic/Latino ^a	474	28.3	198	18.7	672	24.6
Native Hawaiian/other Pacific Islander	2	0.1	1	0.1	3	0.1
White	455	27.2	360	34.0	815	29.8
Multiple races	27	1.6	31	2.9	58	2.1
Total^b	1,672	100	1,060	100	2,732	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.


^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Rates of Diagnoses of HIV Infection among Adults and Adolescents, 2014—United States and 6 Dependent Areas

N = 44,609


Total Rate = 16.6



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Diagnoses of HIV Infection among Adults and Adolescents, by Sex and Race/Ethnicity, 2014—United States



Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.



Diagnoses of HIV Infection among Adult and Adolescent Males, by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	170	18.3
Asian	898	13.4
Black/African American	14,305	94.0
Hispanic/Latino ^a	8,832	41.5
Native Hawaiian/other Pacific Islander	49	22.0
White	10,513	12.6
Multiple races	803	40.6
Total^b	35,571	27.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Diagnoses of HIV Infection among Adult and Adolescent Females, by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	49	5.1
Asian	137	1.8
Black/African American	5,128	30.0
Hispanic/Latino ^a	1,350	6.5
Native Hawaiian/other Pacific Islander	8	3.5
White	1,483	1.7
Multiple races	174	8.2
Total^b	8,328	6.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Hispanics/Latinos can be of any race.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Deaths of Persons with Diagnosed HIV Infection, by Race/Ethnicity, 2013—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	66	2.8
Asian ^a	64	0.4
Black/African American	7,581	19.4
Hispanic/Latino ^b	2,664	4.9
Native Hawaiian/other Pacific Islander	5	0.9
White	5,028	2.5
Multiple races	874	14.1
Total^c	16,281	5.1

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with a diagnosed HIV infection may be due to any cause. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.


^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.




Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Race/Ethnicity, Year-end 2013— United States and 6 Dependent Areas

Males
N = 718,300



Females
N = 232,511



■ American Indian/Alaska Native
■ Asian^a
■ Black/African American

■ Hispanic/Latino^b
■ Native Hawaiian/other Pacific Islander
■ White

■ Multiple races

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.




Adults and Adolescents Living with Diagnosed HIV Infection, by Sex and Transmission Category, Year-end 2013— United States and 6 Dependent Areas

Males
N = 718,300



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU

Females
N = 232,511



- Heterosexual contact^a
- Perinatal exposure
- Other^b

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.


^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.



Rates of Adults and Adolescents Living with Diagnosed HIV Infection, Year-end 2013—United States and 6 Dependent Areas

N = 950,811 Total Rate = 355.9




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Children Aged <13 Years Living with Diagnosed HIV Infection, Year-end 2013—United States and 6 Dependent Areas


N = 2,434

Total Rate = 4.6




Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Stage 3 (AIDS) Classifications and Deaths of Persons with HIV Infection Ever Classified as Stage 3 (AIDS), among Adults and Adolescents, 1985–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting. Deaths of persons with HIV infection, stage 3 (AIDS) may be due to any cause.

Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Race/Ethnicity and Year of Diagnosis, 1985–2014—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with Diagnosed HIV Infection, by Race/Ethnicity and Year of Diagnosis, 1985–2014 – United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Transmission Category and Year of Diagnosis, 1985–2014—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Percentages of Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Transmission Category and Year of Diagnosis, 1985–2014—United States and 6 Dependent Areas




Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Sex and Transmission Category, 2014—United States and 6 Dependent Areas



- Male-to-male sexual contact
- Injection drug use (IDU)
- Male-to-male sexual contact and IDU



- Heterosexual contact^a
- Other^b

Note. All displayed data have been statistically adjusted to account for reporting delays and missing transmission category, but not for incomplete reporting.

^a Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^b Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Stage 3 (AIDS) Classifications among Adults and Adolescents with HIV Infection, by Race/Ethnicity, 2014—United States

Race/ethnicity	No.	Rate
American Indian/Alaska Native	94	4.9
Asian ^a	351	2.5
Black/African American	9,982	30.9
Hispanic/Latino ^b	4,267	10.1
Native Hawaiian/other Pacific Islander	19	4.3
White	5,297	3.1
Multiple races	782	19.0
Total^c	20,792	7.8

Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.

Rates are per 100,000 population.

^a Includes Asian/Pacific Islander legacy cases.

^b Hispanics/Latinos can be of any race.


^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.



Rates of Stage 3 (AIDS) Classifications among Persons with HIV Infection, 2014—United States and 6 Dependent Areas


N = 21,318

Total Rate = 6.6



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Sex, 1993–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), by Race/Ethnicity, 1993–2013—United States and 6 Dependent Areas



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.


^a Hispanics/Latinos can be of any race.

^b Includes Asian/Pacific Islander legacy cases.



Rates of Adults and Adolescents Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2013—United States and 6 Dependent Areas

N = 527,170 Total Rate = 197.4




Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.



Rates of Children Aged <13 Years Living with Diagnosed HIV Infection Ever Classified as Stage 3 (AIDS), Year-end 2013—United States and 6 Dependent Areas

N = 275

Total Rate = 0.5



Note. All displayed data have been statistically adjusted to account for reporting delays, but not for incomplete reporting.